

## **Product datasheet for TA306271**

## **TLR6 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB: 0.5 - 2 ug/mL, ICC: 10 ug/mL

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: TLR6 antibody was raised against a peptide corresponding to 15 amino acids near the amino

terminus of human TLR6.

**Formulation:** PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

**Purification:** Affinity chromatography purified via peptide column

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** toll like receptor 6

Database Link: NP 006059

Entrez Gene 21899 MouseEntrez Gene 10333 Human

Q9Y2C9



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in Drosophila. These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses (1,2). The TLRs act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors (3) so the organism can respond to potential infection. TLR6 was first identified as a close homolog of TLR1, sharing 69% sequence identify (4). Like TLR1, TLR6 can form heterodimers with TLR2, and these TLR6:TLR2 dimers coordinate macrophage activation by Gram-positive bacteria and the yeast cell wall particle zymosan (5). Activation of these complexes not only initiates pro-inflammatory cascades, but also can lead to apoptotic responses (6).

Synonyms: CD286

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Toll-like receptor signaling pathway

## **Product images:**



Western blot analysis of TLR6 in Jurkat cell lysate with TLR6 antibody at (A) 0.5, (B) 1 and (C) 2 ug/mL.

Immunocytochemistry of TLR6 in Jurkat cells with TLR6 antibody at 10 ug/mL.