

Product datasheet for TA306317

IL27 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB: 2 - 4 ug/mL Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: IL-27 antibody was raised against a 14 amino acid peptide from near the center of human IL-

27.

Formulation: PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

Purification: Affinity chromatography purified via peptide column

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: interleukin 27

Database Link: NP 663634

Entrez Gene 246779 MouseEntrez Gene 246778 Human

Q8NEV9

Background: Like interleukin-23 (IL-23), IL-27 is a recently discovered member of the IL-6/IL-12 family of

proinflammatory and immunoregulatory cytokines. It exists as a heterodimer composed of the p40-related protein EBI3 and an IL-12 p35-related protein termed p28. IL-27 is produced

after activation by antigen-presenting cells and induces proliferation of naïve but not

memory CD4+ T-cells. It acts by binding to its receptor WSX-1 and gp130 which results in the activation of a Jak/STAT signaling cascade, suggesting the IL-27 is involved in the regulation of immune processes. It has been suggested that IL-27 can also be used as a therapeutic agent against cancer as it can also induce tumor-specific anti-tumor activity mediated through

CD8+ T-cells, IFN-gamma, and T-bet.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: IL-27; IL-27A; IL27p28; IL30; p28

Protein Families: Druggable Genome, Transmembrane

Product images:



Western blot analysis of IL-27 in Daudi lysate with IL-27 antibody at (A) 2 and (B) 4 μ ml.