

Product datasheet for **TA306319**

AIM (CD5L) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 1 - 4 ug/mL, ICC: 2 ug/mL, IF: 20 ug/mL
Reactivity:	Human, Mouse
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	AIM antibody was raised against a 16 amino acid peptide near the amino terminus of human AIM.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	CD5 molecule like
Database Link:	AAD01446 Entrez Gene 922 Human O43866


Background: Apoptosis inhibitor of macrophages (AIM) is a member of the scavenger receptor cysteine-rich domain superfamily (SRCR-SF) initially identified as an inducible cell surface ligand of CD5. It was shown that AIM functions in the thymus as the inducer of resistance to apoptosis within CD4+/CD8+ thymocytes and as the supporter of the viability of these cells before thymic selection. AIM was also shown to support macrophage survival and enhance their phagocytic function. More recent experiments using recombinant AIM significantly inhibited apoptosis of NKT and T cells obtained from *C. parvum*-stimulated livers in vitro, suggesting that AIM functions to induce resistance to apoptosis in these cells and supports host defense against inflammation during infection.




[View online »](#)

Synonyms: AIM; API6; PRO229; SP-ALPHA; Spalpha


Product images:



Western blot analysis of AIM in human lymph node tissue lysate with AIM antibody at (A) 1, (B) 2 and (C) 4 µg/ml.



Immunohistochemistry of AIM in human lymph node tissue with AIM antibody at 2 µg/ml.



Immunofluorescence of AIM in Human Lymph Node cells with AIM antibody at 20 µg/mL.