

Product datasheet for TA306404

PD1 (PDCD1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1 ug/mL, ICC: 5 ug/mL

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: PD-1 antibody was raised against a 16 amino acid peptide from near the center of human PD-

1.

Formulation: PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

Purification: Affinity chromatography purified via peptide column

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: programmed cell death 1

Database Link: NP 005009

Entrez Gene 18566 MouseEntrez Gene 301626 RatEntrez Gene 5133 Human

Q15116



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


Cell-mediated immune responses are initiated by T lymphocytes that are themselves stimulated by cognate peptides bound to MHC molecules on antig en-presenting cells (APC). T-cell activation is generally self-limited as activated T cells express receptors such as PD-1 (also known as PDCD-1) that mediate inhibitory signals from the APC. PD-1 can bind two different but related ligands, PDL-1 and PDL-2. Upon binding to either of these ligands, signals generated by PD-1 inhibit the activation of the immune response in the absence of "danger signals" such as LPS or other molecules associated with bacteria or other pathogens. Evidence for this is seen in PD1-null mice who exhibit hyperactivated immune systems and autoimmune diseases. Despite its predicted molecular weight, PD-1 often migrates at higher molecular weight in SDS-PAGE.

Synonyms: CD279; hPD-1; hPD-l; hSLE1; PD-1; PD1; SLEB2


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), T cell receptor signaling pathway

Product images:



Western blot analysis of PD-1 in THP-1 cell lysate with PD-1 antibody at 1 ug/ml in the (A) absence and (B) presence of blocking peptide.



Immunohistochemistry of PDCD1 in human brain tissue with PDCD1 antibody at 5 ug/ml.