

## **Product datasheet for TA306481**

## DHX58 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB: 0.5 - 1 ug/mL Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** LGP2 antibody was raised against a 14 amino acid peptide from near the amino terminus of

human LGP2.

**Formulation:** PBS containing 0.02% sodium azide.

**Concentration:** 1ug/ul

**Purification:** Affinity chromatography purified via peptide column

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** DEXH-box helicase 58

Database Link: NP 077024

Entrez Gene 80861 MouseEntrez Gene 303538 RatEntrez Gene 79132 Human

Q96C10

**Background:** Anti-viral innate immune responses are triggered by pathogen-associated molecular patters

(such as the accumulation of intracellular nucleic acids resulting from virus infections) and represent the first line of defense against numerous infectious organisms. The Toll-like receptors (TLR) 3, 7, 8, and 9 are expressed in immune cells and function as transmembrane pattern recognition receptors to detect foreign nucleic acids. Other proteins that play similar roles, such as RIG-1 and MDA5 are members of a CARD-helicase family and are expressed in the cytoplasm. A third protein, LGP2, is similar to RIG-1 and MDA5, except for lacking a

homologous CARD domain. It is thought to act as an element of negative-feedback regulation

of intracellular antiviral signaling. At least four isoforms of LGP2 are known to exist.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: D11LGP2; D11lgp2e; LGP2; RLR-3

**Protein Pathways:** RIG-I-like receptor signaling pathway

## **Product images:**



Western blot analysis of LGP2 in MDA-MB-361 cell lysate with LGP2 antibody at (A) 0.5 and (B) 1 ug/ml.