

Product datasheet for **TA306495**

GRIK1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	WB: 0.5 - 1 ug/mL, ICC: 2.5 ug/mL, IF: 20 ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Grik1 antibody was raised against a 16 amino acid peptide near the center of the human Grik1.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	glutamate ionotropic receptor kainate type subunit 1
Database Link:	NP_000821 Entrez Gene 14805 Mouse Entrez Gene 29559 Rat Entrez Gene 2897 Human P39086



[View online »](#)

Background:

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. Grik1, also known as glutamate receptor 5, belongs to the kainate family of glutamate receptors, which are composed of four subunits and function as ligand-activated ion channels. Grik1 is expressed in GABAergic interneurons of the hippocampus and are thought to participate in the formation of various subtypes of kainate receptors with Grik2 and KA2. Stimulation of Grik1 leads to intracellular calcium release and activation of protein kinase C. Excessive activation has been associated with psychiatric, neurological and neurodegenerative diseases. Numerous isoforms of Grik1 are known to exist and may be subject to RNA editing within the second transmembrane domain, which is thought to alter the properties of ion flow.

Synonyms:


EAA3; EEA3; GLR5; GluK1; GLUR5

Protein Families:


Druggable Genome, Ion Channels: Glutamate Receptors, Transmembrane

Protein Pathways:


Neuroactive ligand-receptor interaction

Product images:

Western blot analysis of Grik1 in rat brain tissue lysate with Grik1 antibody at (A) 0.5 and (B) 1 ug/ml.



Immunohistochemical staining of human brain tissue using Grik1 antibody at 2.5 ug/ml.



Immunofluorescence of Grik1 in Human Brain cells with Grik1 antibody at 20 ug/mL.