

Product datasheet for **TA306576**

SLC35D1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	WB: 1 - 2 ug/mL, ICC: 5 ug/mL
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Slc35D1 antibody was raised against a 20 amino acid peptide near the carboxy terminus of the human Slc35D1.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	solute carrier family 35 member D1
Database Link:	NP_055954 Entrez Gene 23169 Human Q9NTN3



[View online »](#)

Background:


The solute carrier family Slc35 consists of at least 17 proteins that act as nucleotide sugar transporters localized to the Golgi apparatus and endoplasmic reticulum. The role of the ER-resident Slc family member Slc35D1 is to transport both UDP-glucuronic acid and UDP-N-acetylgalactosamine. These molecules can serve as substrates for chondroitin sulfate biosynthesis and mice lacking the Slc35D1 gene developed a lethal form of skeletal dysplasia with severe shortening of limbs and facial structures. Examination of epiphyseal cartilage in these mice revealed a decreased proliferating zone with round chondrocytes, scarce matrices, and reduced proteoglycan aggregates. Loss of function mutations in human Slc35D1 cause Schneckenbecken dysplasia, a severe skeletal dysplasia. This antibody is predicted to not cross-react with the highly homologous Slc35D2.

Synonyms:


UGTREL7

Protein Families:

Transmembrane

Product images:

Western blot analysis of Slc35D1 in Daudi lysate with Slc35D1 antibody at (A) 1 and (B) 2 ug/mL.



Immunocytochemistry of Slc35D1 in Daudi cells with Slc35D1 antibody at 5 ug/mL.