

# Product datasheet for TA306637

## Integrin alpha 4 (ITGA4) Rabbit Polyclonal Antibody

### **Product data:**

#### Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1 ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Integrin alpha 4 antibody was raised against a 13 amino acid peptide from near the carboxy terminus of human Integrin alpha 4.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	integrin subunit alpha 4
Database Link:	<u>NP_000876</u> Entrez Gene 16401 MouseEntrez Gene 311144 RatEntrez Gene 3676 Human <u>P13612</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Integrin alpha 4 (ITGA4) Rabbit Polyclonal Antibody – TA306637
Background:	The integrin alpha 4 (also known as CD49d and ITGA4) belongs to the integrin alpha chain family of proteins. Integrins are heterodimeric integral membrane proteins composed of an alpha and beta chains. Alpha 4 chain associates with either beta 1 or beta 7 chain. It has been demonstrated that the putative ligand-binding sites of both integrin alpha 4 beta 1 and alpha 4 beta 7 is located on the alpha 4 chain. These ligands included Madcam, VCAM, and fibronectin. Madcam is known as the principal ligand for integrin alpha 4 beta 7. Recently it was also demonstrated that HIV-1 envelope can mimic Madcam by binding to and signaling though integrin alpha 4 beta 7, the gut mucosal homing receptor for peripheral T cells. Despite its predicted molecular weight, Integrin alpha 4 will often run at higher molecular weight in SDS-PAGE. At least two isoforms are known to exist. The immunogen to this antibody is outside of the Integrin alpha 4 recombinant protein fragment and thus will not recognize it.
Synonyms:	CD49D; IA4
Protein Families:	Druggable Genome, ES Cell Differentiation/IPS, Transmembrane
Protein Pathway	Arrhythmogenic right ventricular cardiomyopathy (ARVC), Cell adhesion molecules (CAMs), Dilated cardiomyopathy, ECM-receptor interaction, Focal adhesion, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), Leukocyte transendothelial migration, Regulation of actin cytoskeleton

# **Product images:**



> Western blot analysis of Integrin alpha 4 in Jurkat cell lysate with Integrin alpha 4 antibody at 1 ug/ml in (A) the absence and (B) the presence of blocking peptide.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US