

Product datasheet for TA306758

NDP Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB: 1 - 2 ug/mL, ICC: 5 ug/mL

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Norrin antibody was raised against an 18 amino acid peptide from near the amino terminus

of human Norrin.

Formulation: PBS containing 0.02% sodium azide.

Concentration: 1ug/ul

Purification: Affinity chromatography purified via peptide column

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: NDP, norrin cystine knot growth factor

Database Link: NP 000257

Entrez Gene 4693 Human

Q00604



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Norrie disease is an X-linked genetic disorder characterized by progressive atrophy of the eyes, mental disturbances and deafness. The gene responsible for this disease was initially identified through positional cloning. Norrin, the gene product, encodes a small secreted, cysteine-rich protein that is thought to act as a ligand for the Wnt-receptor/b-catenin signal pathway despite having sequence homology with the Wnt family of proteins. Mice lacking this gene have abnormal blood vessel growth in the vitreous and a disorganized retina; transgenic ectopic expression of Norrin restores normal retinal vasculature. Recent evidence shows that Norrin can attenuate tPA and uPA-mediated death of transformed rat retinal ganglion cells (RGC-5) by activating the Wnt/b-catenin pathway and regulating the phosphorylation of LRP-1, a cell surface receptor for tPA and uPA, suggesting the Norrin may function in vivo by regulating kinases which may alter the phosphorylation of LRP-1.


Synonyms: EVR2; FEVR; ND

Protein Families: Druggable Genome, Secreted Protein

Product images:



Western blot analysis of Norrin in Jurkat cell lysate with Norrin antibody at (A) 1 and (B) 2 ug/mL.



Immunocytochemistry of Norrin in Jurkat cells with Norrin antibody at 5 ug/mL.