

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA306793

## **UCMA Rabbit Polyclonal Antibody**

### **Product data:**

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 2.5 - 5 ug/mL
Reactivity:	Human, Mouse
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	UCMA antibody was raised against a 16 amino acid peptide near the carboxy terminus of human UCMA.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	upper zone of growth plate and cartilage matrix associated
Database Link:	<u>NP_660357</u> <u>Entrez Gene 68527 MouseEntrez Gene 221044 Human</u> <u>Q8WVF2</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **CRIGENE** UCMA Rabbit Polyclonal Antibody – TA306793

Background: UCMA is a secreted cartilage-specific protein that was discovered in a screen for differentially expressed genes in retinoic acid-treated mouse chondrocytes. It was also identified in a human chondrocyte EST screen for candidate genes of skeletal dysplasias. UCMA expression is thought to parallel that of collagen II with its expression decreasing with maturation chrondrocytes mature. UCMA is processed by a furin-like protease into two fragments, an amino-terminal fragment and a carboxy-terminal fragment (UCMA-C). Application of recombinant UCMA-C to primary osteoblasts, mesenchymal stem cells, and MC3T3-E1 preosteoblasts interferes with their osteogenic differentiation, but does not affect expression of chondrocyte-specific genes or chondrocyte proliferation, suggesting that UCMA may be involved in the negative control of osteogenic differentiation of osteochondrogenic precursor cells. At least two isoforms of UCMA are known to exist.

Protein Families:

**Product images:** 



Secreted Protein

Western blot analysis of UCMA in SW1353 cell lysate with UCMA antibody at (A) 2.5 and (B) 5 ug/ml.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US