

Product datasheet for **TA306812**

MAK10 (NAA35) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	WB: 1 ug/mL, ICC: 2.5 ug/mL
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	MAK10 antibody was raised against a 19 amino acid peptide near the amino terminus of human MAK10.
Formulation:	PBS containing 0.02% sodium azide.
Concentration:	1ug/ul
Purification:	Affinity chromatography purified via peptide column
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	N(alpha)-acetyltransferase 35, NatC auxiliary subunit
Database Link:	NP_078911 Entrez Gene 64472 Rat Entrez Gene 78689 Mouse Entrez Gene 60560 Human Q5VZE5




[View online »](#)

Background:


The MAK10 gene encodes a 733-amino acid protein with several regions of similarity to T cell receptor alpha-subunit V (variable) regions in yeast. The mammalian homologue of yeast MAK10, also known as EGAP, is one subunit of a novel N-terminal acetyltransferase (NAT) that is highly conserved among vertebrate species. It is expressed in a variety of tissues in the developing rat embryo but restricted in expression in the adult, remaining detectable only in tissues undergoing continual cell renewal or in cells responding to pathological injury. The MAK10-NAT complex is an essential regulatory enzyme controlling the function of a subset of proteins required for embryonic growth control and vessel development. This complex functionally co-assembles in mammalian cells to regulate cell proliferation and is essential for embryonic development, at least in part through the regulation of target of rapamycin (TOR) signaling events. At least two isoforms of MAK10 are known to exist.

Synonyms:

bA379P1.1; EGAP; MAK10; MAK10P

Product images:

Western blot analysis of MAK10 in rat heart tissue lysate with MAK10 antibody at 1 ug/ml in the (A) absence and (B) presence of blocking peptide.



Immunohistochemistry of MAK10 in human heart tissue with MAK10 antibody at 2.5 ug/ml.