

Product datasheet for TA308903

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hydroxysteroid (17 beta) Dehydrogenase 4 (HSD17B4) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: ICC/IF:1:100-1:1000; IHC:1:100-1:1000; WB:1:500-1:3000

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fragment corresponding to a region within amino acids 355 and 645 of

HSD17B4 (Uniprot ID#P51659)

Formulation: 0.1M Tris, 0.1M Glycine, 10% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.

Concentration: lot specific

Purification: Purified by antigen-affinity chromatography.

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80 kDa

Gene Name: hydroxysteroid (17-beta) dehydrogenase 4

Database Link: NP 000405

Entrez Gene 15488 MouseEntrez Gene 3295 Human

P51659

Background: The protein encoded by this gene is a bifunctional enzyme that is involved in the peroxisomal

beta-oxidation pathway for fatty acids. It also acts as a catalyst for the formation of 3-ketoacyl-CoA intermediates from both straight-chain and 2-methyl-branched-chain fatty acids. Defects in this gene that affect the peroxisomal fatty acid beta-oxidation activity are a cause of D-bifunctional protein deficiency (DBPD). An apparent pseudogene of this gene is

present on chromosome 8. [provided by RefSeq]

Synonyms: DBP; MFE-2; MPF-2; PRLTS1; SDR8C1







Protein Families: Druggable Genome

Protein Pathways: Metabolic pathways, Primary bile acid biosynthesis

Product images:




HSD17B4 antibody detects HSD17B4 protein by Western blot analysis. A. 20 ug Rat heart lysate/extract. 7.5% SDS-PAGE. HSD17B4 antibody (TA308903) dilution: 1:3000.



Immunohistochemical analysis of paraffinembedded DLD-1 xenograft, using HSD17B4 (TA308903) antibody at 1:500 dilution.





Immunofluorescence analysis of methanol-fixed Hep3B, using HSD17B4 (TA308903) antibody at 1:500 dilution.