

## **Product datasheet for TA319284**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **IL2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:10,000-1:50,000, WB: 1:1,000-1:5,000, IHC: 1:500-1:2,500

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** This purified antibody was prepared from whole rabbit serum produced by repeated

immunizations with full length recombinant human IL-2 protein.

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Conjugation:** Peroxidase

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: interleukin 2

Database Link: NP 000577

Entrez Gene 3558 Human

P60568

Synonyms: IL-2; lymphokine; TCGF

**Note:** IL-2 is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The

receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in the immune

response to antigenic stimuli.

**Protein Families:** Druggable Genome, Secreted Protein






**Protein Pathways:** 

Allograft rejection, Autoimmune thyroid disease, Cytokine-cytokine receptor interaction, Graft-versus-host disease, Jak-STAT signaling pathway, T cell receptor signaling pathway, Type I diabetes mellitus

## **Product images:**



Western Blot of Rabbit anti-IL-2 Antibody Peroxidase Conjugated Lane 1: Human IL-2 Recombinant Protein Load: 50 ng per lane Secondary antibody: Peroxidase Conjugated IL-2 Antibody at 1:1000 for 60 min at RT Block: MB-070 for 30 min at RT Predicted/Observed size: 16 kDa, 16 kDa