

Product datasheet for TA319287

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL4 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:10,000-1:50,000, WB: 1:1,000-1:5,000, IHC: 1:500;1:2,500

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: This purified antibody was prepared from whole rabbit serum produced by repeated

immunizations with full length recombinant human IL-4 protein.

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: lot specific
Conjugation: Peroxidase

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: interleukin 4

Database Link: NP 000580

Entrez Gene 3565 Human

P05112

Synonyms: BCGF-1; BCGF1; BSF-1; BSF1; IL-4

Note: IL-4 is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for

interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over

120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this

gene encoding distinct isoforms have been reported.






Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Allograft rejection, Asthma, Autoimmune thyroid disease, Cytokine-cytokine receptor

interaction, Fc epsilon RI signaling pathway, Hematopoietic cell lineage, Jak-STAT signaling

pathway, T cell receptor signaling pathway

Product images:



Western Blot of Rabbit Anti - IL-4 Antibody Peroxidase Conjugated Lane 1: Human IL-4 Load: 50 ng per lane Secondary antibody: IL-4 Antibody Peroxidase Conjugated at 1:1,000 for 60 min at RT Block: MB-070 for 30 min RT

Predicted/Observed size: 18 kDa, 18 kDa