

## **Product datasheet for TA319312**

## **IL9 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: ELISA: 1:10,000, WB: 1:1,000, IHC: User Optimized

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** This purified antibody was prepared from whole rabbit serum produced by repeated

immunizations with full length recombinant human IL-9 protein.

**Formulation:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Concentration: lot specific

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** interleukin 9

Database Link: NP 000581

Entrez Gene 3578 Human

P15248

Synonyms: HP40; IL-9; P40

**Note:** IL-9 is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine

stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness. Anti-IL-9 antibody is ideal for investigators involved in Cancer

and Immunology research.

**Protein Families:** Druggable Genome, Secreted Protein



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Protein Pathways:** Asthma, Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

## **Product images:**



anti-Human IL-9 antibody shows detection of a band ~15 kDa in size corresponding to recombinant human IL-9. The identity of the faint higher molecular weight band may represent a homodimer. Molecular weight markers are also shown (left). After transfer, the membrane was blocked overnight with 3% BSA in TBS followed by reaction with primary antibody at a 1:1,000 dilution. Detection occurred using peroxidase conjugated anti-Rabbit IgG (p/n 611-103-122) secondary antibody diluted 1:40,000.