

# **Product datasheet for TA319354**

## **Bach2 Rabbit Polyclonal Antibody**

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1:5000, WB: 1 ug/mL, IP: 1 ug/mL
Reactivity:	Mouse
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Anti-Bach2 antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to the n-terminus region of the Bach2 protein.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	BTB and CNC homology, basic leucine zipper transcription factor 2
Database Link:	<u>NP_031547</u> <u>Entrez Gene 12014 Mouse</u> <u>P97303</u>
Synonyms:	OTTHUMP00000016868



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** Bach2 Rabbit Polyclonal Antibody – TA319354

#### Note:

During an immune response to infectious diseases and cancer, B-cells undergo activation in response to antigen, and differentiate either into short-lived plasma cells, whose numbers contract following withdrawal of the antigenic stimulus, or into memory B-cells. The formation of B-cell memory is the major goal of vaccination, since memory B-cells are able to produce antibodies specific to infections and cancer that can last for the lifetime of the host. Recently it has been shown that the formation of long-lived memory B-cells is dependent upon the transcription factor Bach2, since knockout mice lacking the gene fail to generate class-switched IgG antibody responses and all B-cells undergo default plasma cell differentiation upon antigenic stimulation. While the mechanism by which Bach2 controls Bcell memory fate has been elucidated (it binds and represses the promoter of the gene that encodes the plasma cell master regulatory transcription factor Blimp-1), the upstream mechanism by which the function of Bach2 is regulated is unknown. There is evidence suggesting that Bach2 is phosphorylated in B-cells following stimulation and suspect that this modification may allow the B-cell signaling apparatus to control Bach2 activity and therefore memory fate decisions. Anti-Bach2 is ideal for researchers interested in Cancer and DNA Damage & Repair research.

### **Product images:**



Western Blot of Rabbit anti-Bach2 antibody. Lane 1: 293T cell lysates overexpressing Bach2-Flag. Lane 2: 293T cell lysates. Load: 20 ug per lane. Primary antibody: Bach-2 antibody at 1:1000 for overnight at 4°C. Secondary antibody: rabbit HRP secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 91.7 kDa, ~130 kDa for Bach2. Other band (s): none.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US