

# Product datasheet for TA319421

# Asap1 Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	IF
Recommended Dilution:	ELISA: 1:2,000 - 1:10,000, WB: 1:500 - 1:2,000
Reactivity:	Human, Chicken, Bovine, Dog, Rat, Chimpanzee
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to amino acids 777-787 of mouse ASAP1 protein (see below).
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Gene Name:	ArfGAP with SH3 domain, ankyrin repeat and PH domain1
Database Link:	<u>NP_034156</u> <u>Entrez Gene 50807 HumanEntrez Gene 314961 RatEntrez Gene 482044 Dog</u> <u>Q9QWY8</u>
Synonyms:	AMAP1; CENTB4; DDEF1; DEF-1; KIAA1249; PAG2; PAP; ZG14P




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### Science Asap1 Rabbit Polyclonal Antibody – TA319421

Note:

This antibody is suitable for Cancer, Immunology and Nuclear Signaling research. ASAP1 (also known as AMAP1, 130-kDa phosphatidylinositol 4,5-biphosphate-dependent ARF1 GTPaseactivating protein, PIP2-dependent ARF1 GAP, ADP-ribosylation factor-directed GTPaseactivating protein 1, ARF GTPase-activating protein 1, Development and differentiationenhancing factor 1, Differentiation-enhancing factor 1, DEF-1) is an Arf-directed GTPase activating protein that is a substrate for the kinases Src and FAK and has been implicated in the regulation of membrane traffic, focal adhesions and invadopodia/podosomes. Phosphorylation of ASAP1 at tyrosine 782 has been found to affect enzymatic and some biological activities, including the function of invadopodia. ASAP1 is expressed in many tissues but is most abundant in the testis, brain, lung and spleen. A heightened expression was seen in the adipose tissue from obese (ob) and diabetic (db) animals. Multiple transcript variants have been reported for this protein.

### **Product images:**



Immunofluorescent microscopy using Affinity Purified anti-ASAP1 antibody shows detection of ASAP1present in mouse NIH3T3 cells transfected with activated Src. Specific staining is not present when antibody is pre-incubated with the immunizing peptide prior to reaction with cells. Personal Communication. Paul Randazzo, NIH, CCR, Bethesda, MD.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US