

Product datasheet for TA320214

Caspase-6 (CASP6) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 0.3-1ug/ml, ELISA: 1:128,000

Reactivity: Human
Host: Goat
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide with sequence C-QTEKLDTNITEVD, from the internal region of the protein sequence

according to NP_001217.2; NP_116787.1

Formulation: Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin

Concentration: lot specific

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 33.3 kDa

Gene Name: caspase 6

Database Link: NP 001217

Entrez Gene 839 Human

P55212



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in two transcript variants that encode different isoforms. [provided by RefSeq, Jul 2008]

Synonyms: MCH2

Protein Families: Druggable Genome, Protease, Stem cell - Pluripotency

Protein Pathways: Apoptosis

Product images:



Anti-CASP6 (0.3ug/ml) staining of Human Colon lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by

chemiluminescence.