

## **Product datasheet for TA320296**

## Cd86 Rat Monoclonal Antibody [Clone ID: GL1]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: GL1
Applications: FC

Recommended Dilution: Flow, IHC, Functional Assay, IP

**Reactivity:** Mouse **Host:** Rat

Clonality: Monoclonal

**Formulation:** Aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Concentration:** lot specific

Purification: Affinity purified
Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: CD86 antigen

Database Link: NP 062261

Entrez Gene 12524 Mouse

P42082

Background: The GL1 monoclonal antibody reacts with mouse CD86, an ~80 kDa surface receptor also

known as B7-2. CD86 & CD80 are members of the B7 family of costimulatory molecules. CD86 is expressed at low level on B cells, macrophages, and dendritic cells and is upregulated on B cells through a variety of surface stimuli including the BCR complex, CD40 and some cytokine receptors. CD86 is also expressed by activated mouse T cells and thioglycolate-elicited

peritoneal cells. In addition to CD80 (B7-1), CD86 is a counter-receptor for the T cell surface molecules CD28 and CD152 (CTLA-4). This interaction plays a critical role in T-B crosstalk, T cell costimulation, autoantibody production and Th2-mediated Ig production. The kinetics of upregulation of CD86 upon stimulation, supports its major contribution during the primary

phase of an immune response.

**Synonyms:** B7-2; B70; BU63; CD28LG2; FUN-1; LAB72; MGC34413




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Staining of 3-day LPS-stimulated BALB/c splenocytes with 0.25 ug of Rat IgG2a kappa Isotype Control Purified (open histogram) or 0.25 ug of Purified Anti-Mouse CD86 (B7-2) (filled histogram) followed by Anti-Rat IgG FITC. Total viable cells were used for analysis.