

### **Product datasheet for TA320470**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TBX21 Mouse Monoclonal Antibody [Clone ID: eBio4B10 (4B10, 4-B10)]

#### **Product data:**

**Product Type:** Primary Antibodies

**Clone Name:** eBio4B10 (4B10, 4-B10)

Applications: WB

Recommended Dilution: IHC, IP, WB

Reactivity: Human, Mouse, Rhesus Monkey

Host: Mouse

Clonality: Monoclonal

**Formulation:** Aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

**Concentration:** lot specific

Purification: Affinity purified
Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: T-box 21

Database Link: NP 037483

Entrez Gene 57765 MouseEntrez Gene 30009 Human

Q9UL17

Background: The eBio4B10 monoclonal antibody reacts with mouse and human T-bet. T-bet is a Th1-

specific T-box transcription factor critical to the development of the Th1 CD4+ T cell lineage. This is known based on the observations that T-Bet deficient mice have impaired Th1 cell development, and that ectopic expression of T-Bet results in development skewed to the Th1 lineage. T-Bet expression is induced by the Th1 cytokine IFN gamma, and T-Bet also regulates

the expression of IFN gamma, likely, at least in part, through the modification of DNA accessibility and histone remodeling. In addition to IFN gamma, T-Bet is also known to regulate the expression of IL-12R beta and IL-2. Moreover, T-Bet plays a role in class-switch

recombination in B-cells.


**Synonyms:** T-bet; T-PET; TBET; TBLYM

**Protein Families:** Druggable Genome, Transcription Factors





# **Product images:**



CD4+ T cells were sorted from mouse spleen, activated with Anti-Mouse CD3 and Anti-Mouse CD28, followed by culture in Th1-polarizing conditions, and re-activation with PMA and lonomycin. Lysates from control (left) or PMA and lonomycin-re-activated (right) cells were probed with Anti-Human/Mouse T-bet Purified at 2 ug/mL, and revealed with Anti-Mouse IgG HRP.