

Product datasheet for TA320480

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TLR2 Mouse Monoclonal Antibody [Clone ID: TL2.1]

Product data:

Product Type: Primary Antibodies

Clone Name: TL2.1
Applications: FC

Recommended Dilution: Flow, IHC, IP, WB

Reactivity: Human Host: Mouse

Clonality: Monoclonal

Formulation: Aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Concentration: lot specific

Purification: Affinity purified
Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: toll like receptor 2

Database Link: NP 003255

Entrez Gene 7097 Human

<u>O60603</u>

Background: The TL2.1 monoclonal antibody reacts with human Toll-like receptor 2 (TLR2). To date, at least


ten members of the Toll family have been identified in human. This family of type I transmembrane proteins is characterized by an extracellular domain with leucine-rich repeats and a cytoplasmic domain with homology to the type I IL-1 receptor. Two of these receptors, TLR2 and TLR4, are pattern recognition receptors and signaling molecules in response to bacterial lipoproteins and have been implicated in innate immunity and inflammation. TLR2 is expressed by peripheral blood monocytes and is responsible for distinguishing different pathogens. TL2.1, a blocking antibody, has been used to study the role of TLR-2 as a pattern recognition receptor in microbial lipoprotein/lipopeptide induced cytokine production from human peripheral blood mononuclear cells. TL2.1 has been reported to immunoprecipitate human TLR2 (~90 kDa) from PBMC and HMEC.



Synonyms: CD282; TIL4

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Toll-like receptor signaling pathway

Product images:



Human peripheral blood monocytes were stained with Anti-Human CD282 (TLR2) Purified followed by Anti-Mouse IgG FITC. (Gated on normal peripheral monocytes)