

## **Product datasheet for TA321038**

## **Angiotensinogen (AGT) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Human liver cancer tissue

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide peptide corresponding to a region derived from 34-41 amino acids of

human angiotensinogen (serpin peptidase inhibitor, clade A, member 8)

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 53 kDa

**Gene Name:** angiotensinogen

Database Link: NP 000020

Entrez Gene 183 Human

P01019



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

The protein encoded by this gene; pre-angiotensinogen or angiotensinogen precursor; is expressed in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. The resulting product; angiotensin I; is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. The protein is involved in maintaining blood pressure and in the pathogenesis of essential hypertension and preeclampsia. Mutations in this gene are associated with susceptibility to essential hypertension; and can cause renal tubular dysgenesis; a severe disorder of renal tubular development. Defects in this gene have also been associated with non-familial structural atrial fibrillation; and inflammatory bowel disease.

Synonyms: ANHU; SERPINA8

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Renin-angiotensin system

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human liver cancer tissue

Primary antibody: TA321038 (AGT Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 5 minutes