

## **Product datasheet for TA321046**

## **ALDOB Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human fetal liver tissue

IHC: 50-200

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 109-122 amino acids of human

aldolase B, fructose-bisphosphate

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 39 kDa

**Gene Name:** aldolase, fructose-bisphosphate B

Database Link: NP 000026

Entrez Gene 24190 RatEntrez Gene 230163 MouseEntrez Gene 229 Human

P05062



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

Fructose-1,6-bisphosphate aldolase (EC 4.1.2.13) is a tetrameric glycolytic enzyme that catalyzes the reversible conversion of fructose-1,6-bisphosphate to glyceraldehyde 3-phosphate and dihydroxyacetone phosphate. Vertebrates have 3 aldolase isozymes which are distinguished by their electrophoretic and catalytic properties. Differences indicate that aldolases A, B, and C are distinct proteins, the products of a family of related 'housekeeping' genes exhibiting developmentally regulated expression of the different isozymes. The developing embryo produces aldolase A, which is produced in even greater amounts in adult muscle where it can be as much as 5% of total cellular protein. In adult liver, kidney and intestine, aldolase A expression is repressed and aldolase B is produced. In brain and other nervous tissue, aldolase A and C are expressed about equally. There is a high degree of homology between aldolase A and C. Defects in ALDOB cause hereditary fructose intolerance.


Synonyms: ALDB; ALDO2

**Protein Families:** Druggable Genome

**Protein Pathways:** Fructose and mannose metabolism, Glycolysis / Gluconeogenesis, Metabolic pathways,

Pentose phosphate pathway

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Human fetal liver tissue

Primary antibody: TA321046 (ALDOB Antibody) at


dilution 1/260

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 10 seconds





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321046 (ALDOB Antibody) at dilution 1/55 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321046 (ALDOB Antibody) at dilution 1/55, treated with synthetic peptide. (Original magnification: ×200)