

### **Product datasheet for TA321113**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# alpha Synuclein (SNCA) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500-1000

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Peptide sequence around aa.123~127 (E-A-Y-E-M ) derived from Human a-Synuclein.

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 14 kDa

**Gene Name:** synuclein alpha

Database Link: NP 000336

Entrez Gene 6622 Human

P37840

**Background:** SncA is a member of the synuclein family of structurally related proteins that are prominently

expressed in the central nervous system, which also includes beta- and gamma-synuclein.

Synucleins are abundantly expressed in the brain and SncA and Snc-Beta inhibit phospholipase D2 selectively. SncA may serve to integrate presynaptic signaling and membrane trafficking. Aggregated SncA proteins form brain lesions that are hallmarks of neurodegenerative synucleinopathies. Defects in SncA play a role in the pathogenesis of Parkinson disease. SncA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer disease. SncA shares 95% sequence homology with rat SncA. Rat SncA is specifically expressed in brain and is associated with synaptosomal membranes in neurons




Synonyms: NACP; PARK1; PARK4; PD1

**Protein Families:** Druggable Genome

**Protein Pathways:** Alzheimer's disease, Parkinson's disease

# **Product images:**



Predicted band size: 14 kDa. Positive control: Mouse Brain tissue lysate. Recommended dilution: 1/500-1000. (Gel: 12%SDS-PAGE Lysate: 30 ug Primary antibody: 1/500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 minute)