

Product datasheet for TA321142S

CYP1A1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse liver tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to C terminal 300 amino acids of human cytochrome P450,

family 1, subfamily A, polypeptide 1

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 58 kDa

Gene Name: cytochrome P450 family 1 subfamily A member 1

Database Link: NP 000490

Entrez Gene 13076 MouseEntrez Gene 24296 RatEntrez Gene 1543 Human

P04798



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CYP1A1 Rabbit Polyclonal Antibody - TA321142S

Background: This gene; CYP1A1; encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol; steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs); some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however; it is able to metabolize some PAHs to

carcinogenic intermediates. The gene has been associated with lung cancer risk. A related family member; CYP1A2; is located approximately 25 kb away from CYP1A1 on chromosome


15.

Synonyms: AHH; AHRR; CP11; CYP1; CYPIA1; P1-450; P450-C; P450DX

Protein Families: Druggable Genome, P450, Transmembrane

Protein Pathways: Metabolism of xenobiotics by cytochrome P450, Retinol metabolism, Tryptophan metabolism

Product images:



Gel: 10%SDS-PAGE Lysate: 30 μg

Lane: Mouse liver tissue

Primary antibody: [TA321142] (CYP1A1 Antibody)


at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 seconds





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321142] (CYP1A1 Antibody) at dilution 1/25 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321142] (CYP1A1 Antibody) at dilution 1/25, treated with fusion protein. (Original magnification: ×200)