

Product datasheet for **TA321146**

Factor X (F10) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 200-1000 WB positive control: A431 cells
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Fusion protein corresponding to a region derived from 235-488 amino acids of human coagulation factor X
Formulation:	PBS pH7.3, 0.05% NaN ₃ , 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	55 kDa
Gene Name:	coagulation factor X
Database Link:	NP_000495 Entrez Gene 14058 Mouse Entrez Gene 29243 Rat Entrez Gene 2159 Human P00742



[View online »](#)

Background:

This gene encodes the vitamin K-dependent coagulation factor X of the blood coagulation cascade. This factor undergoes multiple processing steps before its preproprotein is converted to a mature two-chain form by the excision of the tripeptide RKR. Two chains of the factor are held together by 1 or more disulfide bonds; the light chain contains 2 EGF-like domains; while the heavy chain contains the catalytic domain which is structurally homologous to those of the other hemostatic serine proteases. The mature factor is activated by the cleavage of the activation peptide by factor IXa (in the intrinsic pathway); or by factor VIIa (in the extrinsic pathway). The activated factor then converts prothrombin to thrombin in the presence of factor Va; Ca²⁺; and phospholipid during blood clotting. Mutations of this gene result in factor X deficiency; a hemorrhagic condition of variable severity.

Synonyms:


FX; FXA

Protein Families:

Druggable Genome, Protease, Transmembrane

Protein Pathways:

Complement and coagulation cascades

Product images:

Gel: 10%SDS-PAGE

Lysate: 40 µg

Lane: A431 cells

Primary antibody: TA321146 (F10 Antibody) at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 10 seconds