

Product datasheet for TA321153

p53 (TP53) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: HT29, HepG2, Hela, 231 cell lysates

Reactivity: Human
Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 1-233 amino acids of human tumor

protein p53

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 44 kDa

Gene Name: tumor protein p53

Database Link: NP 000537

Entrez Gene 7157 Human

P04637



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

This gene encodes tumor protein p53; which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest; apoptosis; senescence; DNA repair; or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines; where it's believed to contribute to transformation and malignancy, p53 is a DNA-binding protein containing transcription activation; DNA-binding; and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion; and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site; and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies; but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms; which can regulate p53 transcriptional activity.

Synonyms: BCC7; LFS1; P53; TRP53


Protein Families: Druggable Genome, Stem cell - Pluripotency, Transcription Factors

Protein Pathways: Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle,

Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer,

Small cell lung cancer, Thyroid cancer, Wnt signaling pathway

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-4: HT29 HepG2 Hela 231 cell lysates Primary antibody: TA321153 (TP53 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at

1/5000 dilution

Exposure time: 30 seconds