

Product datasheet for TA321166S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Alpha 1 Acid Glycoprotein (ORM1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Human testis and prostate tissue

IHC: 15-50

Positive control: Human lung cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 19-201 amino acids of human

orosomucoid 1

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 24 kDa

Gene Name: orosomucoid 1 **Database Link:** NP 000598

Entrez Gene 5004 Human

P02763

Background: This gene encodes a key acute phase plasma protein. Because of its increase due to acute

inflammation, this protein is classified as an acute-phase reactant. The specific function of

this protein has not yet been determined; however, it may be involved in aspects of

immunosuppression.


Synonyms: AGP-A; AGP1; HEL-S-153w; ORM






Protein Families: Druggable Genome, Secreted Protein

Product images:






Gel: 12%SDS-PAGE Lysate: 40 µg Lane 1-2: Human testis tissue Human prostate tissue Primary antibody: [TA321166] (ORM1 Antibody) at dilution 1/160 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 2 minutes

Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA321166] (ORM1 Antibody) at dilution 1/15 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human lung cancer tissue using [TA321166] (ORM1 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)