

## **Product datasheet for TA321202S**

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## Dopamine Receptor D4 (DRD4) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-1000

WB positive control: Mouse skeletal muscle tissue

IHC: 5-20

Positive control: Human cervical cancer

Predicted cell location: Cytoplasm, Cell membrane

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 2-15 amino acids of Human

Dopamine receptor D4

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 48 kDa

**Gene Name:** dopamine receptor D4

Database Link: NP 000788

Entrez Gene 13491 MouseEntrez Gene 1815 Human

P21917





**Background:** This gene encodes the D4 subtype of the dopamine receptor. The D4 subtype is a G-protein

coupled receptor which inhibits adenylyl cyclase. It is a target for drugs which treat schizophrenia and Parkinson disease. Mutations in this gene have been associated with various behavioral phenotypes; including autonomic nervous system dysfunction; attention deficit/hyperactivity disorder; and the personality trait of novelty seeking. This gene contains a polymorphic number (2-10 copies) of tandem 48 nt repeats; the sequence shown contains


four repeats.

Synonyms: D4DR

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 30 μg

Lane: Mouse skeletal muscle tissue


Primary antibody: [TA321202] (DRD4 Antibody) at

dilution 1/350

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321202] (DRD4 Antibody) at dilution 1/5 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321202] (DRD4 Antibody) at dilution 1/5, treated with synthetic peptide. (Original magnification: ×200)