

Product datasheet for TA321212

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Metabotropic Glutamate Receptor 3 (GRM3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human brain

Predicted cell location: Cytoplasm, Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 23-276 amino acids of human

glutamate receptor, metabotropic 3

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: glutamate metabotropic receptor 3

Database Link: NP 000831

Entrez Gene 24416 RatEntrez Gene 108069 MouseEntrez Gene 2913 Human

Q14832





Background:


L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors; that have been divided into 3 groups on the basis of sequence homology; putative signal transduction mechanisms; and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4; GRM6; GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities.?

Synonyms: GLUR3; GPRC1C; mGlu3; MGLUR3

Protein Families: Druggable Genome, GPCR, Transmembrane


Protein Pathways: Neuroactive ligand-receptor interaction

Product images:




Immunohistochemistry of paraffin-embedded Human brain tissue using TA321212 (GRM3 Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human brain tissue using TA321212 (GRM3 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321212 (GRM3 Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA321212 (GRM3 Antibody) at dilution 1/50, treated with fusion protein. (Original magnification: ×200)