

Product datasheet for TA321213S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Metabotropic Glutamate Receptor 3 (GRM3) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: 293T cells

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 366-380 amino acids of Human

glutamate receptor, metabotropic 3

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 99 kDa

Gene Name: glutamate metabotropic receptor 3

Database Link: NP 000831

Entrez Gene 24416 RatEntrez Gene 108069 MouseEntrez Gene 2913 Human

Q14832



Metabotropic Glutamate Receptor 3 (GRM3) Rabbit Polyclonal Antibody - TA321213S

Background: L-glutamate is the major excitatory neurotransmitter in the central nervous system and

AMP cascade but differ in their agonist selectivities.

activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors; that have been divided into 3 groups on the basis of sequence homology; putative signal transduction mechanisms; and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4; GRM6; GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic

Synonyms: GLUR3; GPRC1C; mGlu3; MGLUR3

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: 293T cells

Primary antibody: [TA321213] (GRM3 Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 2 minutes