

Product datasheet for TA321229S

CD10 (MME) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 50-200

Positive control: Human brain Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide peptide corresponding to a region derived from 519-533 amino acids of

human membrane metallo-endopeptidase

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: membrane metallo-endopeptidase

Database Link: NP 000893

Entrez Gene 17380 MouseEntrez Gene 24590 RatEntrez Gene 4311 Human

P08473



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype; which represent 85% of cases of ALL. This protein is not restricted to leukemic cells; however; and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney; where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon; enkephalins; substance P; neurotensin; oxytocin; and bradykinin. This gene; which encodes a 100-kD type II transmembrane glycoprotein; exists in a single copy of greater than 45 kb. The 5' untranslated region of this gene is alternatively spliced; resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing.?

Synonyms: CALLA; CD10; CMT2T; NEP; SCA43; SFE

Protein Families: Druggable Genome, Protease, Transmembrane


Protein Pathways: Alzheimer's disease, Hematopoietic cell lineage, Renin-angiotensin system

Product images:



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA321229] (MME Antibody) at dilution 1/85 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human brain tissue using [TA321229] (MME Antibody) at dilution 1/85, treated with synthetic peptide. (Original magnification: ×200)