

Product datasheet for TA321241S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Prostaglandin D Synthase (PTGDS) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela cell, Human muscle tissue lysates

IHC: 50-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 64-325 amino acids of human

prostaglandin D2 synthase 21kDa (brain)prostaglandin D2 synthase 21kDa (brain)

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21 kDa

Gene Name: prostaglandin D2 synthase

Database Link: NP 000945

Entrez Gene 19215 MouseEntrez Gene 5730 Human

P41222



Prostaglandin D Synthase (PTGDS) Rabbit Polyclonal Antibody - TA321241S


Background:

The protein encoded by this gene is a glutathione-independent prostaglandin D synthase that catalyzes the conversion of prostaglandin H2 (PGH2) to postaglandin D2 (PGD2). PGD2 functions as a neuromodulator as well as a trophic factor in the central nervous system. PGD2 is also involved in smooth muscle contraction/relaxation and is a potent inhibitor of platelet aggregation. This gene is preferentially expressed in brain. Studies with transgenic mice overexpressing this gene suggest that this gene may be also involved in the regulation of non-rapid eye movement sleep.


Synonyms: L-PGDS; LPGDS; PGD2; PGDS; PGDS2

Protein Pathways: Arachidonic acid metabolism, Metabolic pathways

Product images:




Gel: 8%SDS-PAGE Lysate: 40 µg Lane 1-2: Hela cell Human muscle tissue lysates Primary antibody: [TA321241] (MAP2K3 Antibody) at dilution 1/400 Secondary antibody: Goat anti rabbit IgG at 1/5000 dilution Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321241] (MAP2K3 Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321241] (MAP2K3 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321241] (MAP2K3 Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321241] (MAP2K3 Antibody) at dilution 1/60, treated with fusion protein. (Original magnification: ×200)