

## **Product datasheet for TA321248S**

## PAF Receptor (PTAFR) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 1-14 amino acids of Human

platelet-activating factor receptor

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** platelet activating factor receptor

Database Link: NP 000943

Entrez Gene 5724 Human

P25105



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes a seven-transmembrane G-protein-coupled receptor for platelet-activating factor (PAF) that localizes to lipid rafts and/or caveolae in the cell membrane. PAF (1-0-alkyl-2-acetyl-sn-glycero-3-phosphorylcholine) is a phospholipid that plays a significant role in oncogenic transformation; tumor growth; angiogenesis; metastasis; and pro-inflammatory processes. Binding of PAF to the PAF-receptor (PAFR) stimulates numerous signal transduction pathways including phospholipase C; D; A2; mitogen-activated protein kinases (MAPKs); and the phosphatidylinositol-calcium second messenger system. Following PAFR activation; cells become rapidly desensitized and this refractory state is dependent on PAFR phosphorylation; internalization; and down-regulation. Alternative splicing results in multiple transcript variants.

Synonyms: PAFR

**Protein Families:** Druggable Genome, GPCR, Transmembrane


**Protein Pathways:** Calcium signaling pathway, Neuroactive ligand-receptor interaction

## **Product images:**



Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA321248] (PTAFR Antibody) at dilution 1/70 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using [TA321248] (PTAFR Antibody) at dilution 1/70, treated with synthetic peptide. (Original magnification: ×200)