

Product datasheet for TA321266

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Somatostatin Receptor 5 (SSTR5) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 500-2000

WB positive control: Human heart tissue lysate

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 320-334 amino acids of human

somatostatin receptor 5

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 39 kDa

Gene Name: somatostatin receptor 5

Database Link: NP 001044

Entrez Gene 6755 Human

P35346



Background:


Somatostatin and its related peptide cortistatin exert multiple biological actions on normal and tumoral tissue targets by interacting with somatostatin receptors (SSTRs). The protein encoded by this gene is one of the SSTRs; which is a multi-pass membrane protein and belongs to the G-protein coupled receptor 1 family. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase; and different regions of this receptor molecule are required for the activation of different signaling pathways. A mutation in this gene results in somatostatin analog resistance. Alternatively spliced transcript variants have been identified in this gene.

Synonyms: SS-5-R

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human heart tissue lysate

Primary antibody: TA321266 (SSTR5 Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 30 seconds