

## **Product datasheet for TA321267**

## **Transferrin (TF) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Human kidney and prostate tissue

IHC: 25-100

Positive control: Human prostate cancer

Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 562-577 amino acids of Human

Transferrin

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 77 kDa

Gene Name: transferrin

Database Link: NP 001054

Database Link: NP 001054

Entrez Gene 7018 Human

P02787



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine; reticuloendothelial system; and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum.

Synonyms: HEL-S-71p; PRO1557; PRO2086; TFQTL1


Protein Families: Druggable Genome, Secreted Protein

## **Product images:**




Gel: 6%SDS-PAGE
Lysate: 40 µg
Lane 1-2: Human kidney tissue
Human prostate tissue
Primary antibody: TA321267 (TF Antibody) at
dilution 1/200
Secondary antibody: Goat anti rabbit IgG at
1/8000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA321267 (TF Antibody) at dilution 1/20 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA321267 (TF Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)