

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA321272S

### Somatostatin Receptor 2 (SSTR2) Rabbit Polyclonal Antibody

### **Product data:**

Applications:WBRecommended Dilution:WB: 200-1000 WB positive control: Mouse thymus tissueReactivity:Human, Mouse, RatHost:RabitHost:BabbitIsotype:IgGClonalty:OhylonalImmunogen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human sonatostatin receptor 2Formulation:PS pH7.3, 0.05% NAN3, 50% glycerolPurification:NonojugatedYongation:UnconjugatedStorage:Stora erceived.Storage:Stole for 12 months from date of receipt.Predicted Protein Size:NP. 001041 Triter Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human PasiZMABackground:Smotostatin acts at many sites to inhibit the release of many hormones and other secretor the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven transmembrane segments and is expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven transmembrane segments and is expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors sharing seven transmembrane segments and is expressed in a tissue-specific manne	Product Type:	Primary Antibodies
IdentifiedIdentifiedReactivity:Human, Mouse, RatHost:RabitIsotype:IgClonality:PolyclonalImmongen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Nition agritForngugation:Nition agritStorage:Stora et -20°C as received.Fordity:Stabilor 12 months from date of receipt.Predicted Protein SizeAlkoGene Name:NP.001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human 20324Fackground:Somatostatin acts at many sites to inhibit the release of frames of spanies	Applications:	WB
Host:RabitIsotype:IgGClonality:PolyclonalImmunogen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Antigen affinity purificationConjugation:UnconjugatedStorage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size14 kDaGene Name:Somatostatin arceptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human 20874Background:Somatostatin atts at many sites to inhibit the release of many hormones and other server of the superfamily of receptors having seven transmembrane segments and is expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having seven transmembrane segments and is expressed in higherst superfamily of receptors having s	Recommended Dilution:	
Isotype:IgGClonality:PolyclonalImmunogen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Natigen affinity purificationConjugation:UnconjugatedStorage:Sorre at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:At kDaGene Name:Somatostatin receptor 2NP. 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human 20874Sakground:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory superfamily of receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highers.Synonyms:SIR-1; SS2R	Reactivity:	Human, Mouse, Rat
Clonality:PolyclonalImmunogen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Antigen affinity purificationConjugation:UnconjugatedStorage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Host:	Rabbit
Immunogen:Synthetic peptide corresponding to a region derived from 355-369 amino acids of human somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Antigen affinity purificationConjugation:UnconjugatedStorage:Stole of a 1-20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:Antigon affinity exploredGene Name:Somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretor by proteins. The biologic effects of somatostatin are probably mediated by a family of g proteins interesting seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having seven transmembrane segments and is expressed in highest is uperfamily of receptors having s	lsotype:	lgG
Formulation:Somatostatin receptor 2Formulation:PBS pH7.3, 0.05% NaN3, 50% glycerolPurification:Antigen affinity purificationConjugation:UnconjugatedStorage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Clonality:	Polyclonal
Purification:Antigen affinity purificationConjugation:UnconjugatedStorage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- soupel receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Immunogen:	
Conjugation:UnconjugatedStorage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SIF-1; SS2R	Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Storage:Store at -20°C as received.Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Purification:	Antigen affinity purification
Stability:Stable for 12 months from date of receipt.Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- superfamily of receptors having seven transmembrane segments and is expressed in highestSynonyms:SRIF-1; SS2R	Conjugation:	Unconjugated
Predicted Protein Size:41 kDaGene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Storage:	Store at -20°C as received.
Gene Name:somatostatin receptor 2Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein superfamily of receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the evels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Stability:	Stable for 12 months from date of receipt.
Database Link:NP 001041 Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Predicted Protein Size:	41 kDa
Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human P30874Background:Somatostatin acts at many sites to inhibit the release of many hormones and other secretory proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Gene Name:	somatostatin receptor 2
proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest levels in cerebrum and kidney.Synonyms:SRIF-1; SS2R	Database Link:	Entrez Gene 20606 MouseEntrez Gene 54305 RatEntrez Gene 6752 Human
	Background:	proteins. The biologic effects of somatostatin are probably mediated by a family of G protein- coupled receptors that are expressed in a tissue-specific manner. SSTR2 is a member of the superfamily of receptors having seven transmembrane segments and is expressed in highest
Protein Families: Druggable Genome, GPCR, Transmembrane	Synonyms:	SRIF-1; SS2R
	Protein Families:	Druggable Genome, GPCR, Transmembrane




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Protein Pathways:

Neuroactive ligand-receptor interaction

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 60 µg Lane: Mouse thymus tissue Primary antibody: [TA321272] (SSTR2 Antibody) at dilution 1/200 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution Exposure time: 1 minute

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US