

## **Product datasheet for TA321306**

## Caspase-6 (CASP6) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** WB

Recommended Dilution: WB: 50-500

WB positive control: Mouse stomach tissue

Reactivity: Human, Mouse, Rat

Host: Rabbit **IgG** Isotype:

Clonality: Polyclonal

Fusion protein corresponding to a region derived from 66-221 amino acids of human caspase Immunogen:

6, apoptosis-related cysteine peptidase

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

**Predicted Protein Size:** 33 kDa Gene Name: caspase 6 Database Link: NP 001217

Entrez Gene 12368 MouseEntrez Gene 83584 RatEntrez Gene 839 Human

P55212

Background: This gene encodes a protein which is a member of the cysteine-aspartic acid protease

(caspase) family. Sequential activation of caspases plays a central role in the execution-phase

of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits; large and small; that dimerize to form the active enzyme. This protein is processed by caspases 7; 8 and 10; and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative

splicing of this gene results in two transcript variants that encode different isoforms.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: MCH2

**Protein Families:** Druggable Genome, Protease, Stem cell - Pluripotency

**Protein Pathways:** Apoptosis

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 μg

Lane: Mouse stomach tissue

Primary antibody: TA321306 (CASP6 Antibody) at

dilution 1/50

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 30 seconds