

Product datasheet for TA321403S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Creatine kinase M type (CKM) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Mouse muscle and Heart tissue

IHC: 50-200

Positive control: Human esophagus cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 14-26 amino acids of Human

creatine kinase, muscle

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43 kDa

Gene Name: creatine kinase, M-type

Database Link: NP 001815

Entrez Gene 12715 MouseEntrez Gene 24265 RatEntrez Gene 1158 Human

P06732





Creatine kinase M type (CKM) Rabbit Polyclonal Antibody - TA321403S

Background: The protein encoded by this gene is a cytoplasmic enzyme involved in energy homeostasis

and is an important serum marker for myocardial infarction. The encoded protein reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in striated muscle as well as in other tissues; and as a heterodimer with a similar brain isozyme in heart. The encoded protein is a member of the


ATP:guanido phosphotransferase protein family

Synonyms: CKMM; M-CK

Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, Metabolic pathways

Product images:



Gel: 10%SDS-PAGE Lysate: 50 μg

Lane 1-2: Mouse muscle tissue

Mouse Heart tissue


Primary antibody: [TA321403] (CKM Antibody) at

dilution 1/950

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 seconds



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA321403] (CKM Antibody) at dilution 1/50 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using [TA321403] (CKM Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)