

## **Product datasheet for TA321411S**

## **DUT Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: 293T cells

IHC: 10-50

Positive control: Human brain Predicted cell location: Cytoplasm

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 132-244 amino acids of Human

Deoxyuridine triphosphatase

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 27 kDa

**Gene Name:** deoxyuridine triphosphatase

Database Link: NP 001939

Entrez Gene 497778 RatEntrez Gene 1854 Human

P33316



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:


This gene encodes an essential enzyme of nucleotide metabolism. The encoded protein forms a ubiquitous; homotetrameric enzyme that hydrolyzes dUTP to dUMP and pyrophosphate. This reaction serves two cellular purposes: providing a precursor (dUMP) for the synthesis of thymine nucleotides needed for DNA replication; and limiting intracellular pools of dUTP. Elevated levels of dUTP lead to increased incorporation of uracil into DNA; which induces extensive excision repair mediated by uracil glycosylase. This repair process; resulting in the removal and reincorporation of dUTP; is self-defeating and leads to DNA fragmentation and cell death. Alternative splicing of this gene leads to different isoforms that localize to either the mitochondrion or nucleus. A related pseudogene is located on chromosome 19.

**Synonyms:** dUTPase

**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, Pyrimidine metabolism

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 40 µg Lane: 293T cells

Primary antibody: [TA321411] (DUT Antibody) at


dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 minutes





Immunohistochemistry of paraffin-embedded Human brain tissue using [TA321411] (DUT Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using [TA321411] (DUT Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)