

## **Product datasheet for TA321469S**

## **LOX Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Rat brain tissue lysate

IHC: 20-100

Positive control: Human liver cancer Predicted cell location: Secreted

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 248-262 amino acids of human

lysyl oxidase

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 47 kDa

Gene Name: lysyl oxidase

Database Link: NP 002308

Entrez Gene 16948 MouseEntrez Gene 24914 RatEntrez Gene 4015 Human

P28300



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The protein encoded by this gene is an extracellular copper enzyme that initiates the crosslinking of collagens and elastin. The enzyme catalyzes oxidative deamination of the epsilon-amino group in certain lysine and hydroxylysine residues of collagens and lysine residues of elastin. In addition to crosslinking extracellular matrix proteins; the encoded protein may have a role in tumor suppression. Defects in this gene are a cause of autosomal recessive cutis laxa type I (CL type I). Two transcript variants encoding different isoforms have been found for this gene.

Synonyms: MGC105112

**Protein Families:** Druggable Genome, Secreted Protein

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Rat brain tissue lysate


Primary antibody: [TA321469] (LOX Antibody) at

dilution 1/200

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 90 seconds



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321469] (LOX Antibody) at dilution 1/20 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321469] (LOX Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)