

### **Product datasheet for TA321496**

## **OGG1 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500-2000

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39 kDa

**Gene Name:** 8-oxoguanine DNA glycosylase

Database Link: NP 002533

Entrez Gene 18294 MouseEntrez Gene 81528 RatEntrez Gene 4968 Human

O15527

**Background:** This gene encodes the enzyme responsible for the excision of 8-oxoguanine, a mutagenic

base byproduct which occurs as a result of exposure to reactive oxygen. The action of this enzyme includes lyase activity for chain cleavage. Alternative splicing of the C-terminal region of this gene classifies splice variants into two major groups, type 1 and type 2, depending on the last exon of the sequence. Type 1 alternative splice variants end with exon 7 and type 2 end with exon 8. All variants share the N-terminal region in common, which contains a mitochondrial targeting signal that is essential for mitochondrial localization. Many alternative splice variants for this gene have been described, but the full-length nature for

every variant has not been determined.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### OGG1 Rabbit Polyclonal Antibody - TA321496

Synonyms: HMMH; HOGG1; MUTM; OGH1

Protein Families: Druggable Genome
Protein Pathways: Base excision repair

# **Product images:**



Predicted band size: 39 kDa. Positive control: Human heart tissue lysate. Recommended dilution: 1/500-2000. (Gel: 10%SDS-PAGE Lysate: 40 ug Primary antibody: 1/500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 1 minute)