

Product datasheet for TA321511

PCNA Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Hela cells

IHC: 50-200

Positive control: Human gasrtic cancer Predicted cell location: Cytoplasm, Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Fusion protein corresponding to a region derived from 9-252 amino acids of Human

Proliferating cell nuclear antigen

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29 kDa

Gene Name: proliferating cell nuclear antigen

Database Link: NP 002583

Entrez Gene 18538 MouseEntrez Gene 25737 RatEntrez Gene 5111 Human

P12004



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

PCNA Rabbit Polyclonal Antibody - TA321511

Background: The protein encoded by this gene is found in the nucleus and is a cofactor of DNA

polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage; this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of

this gene have been described on chromosome 4 and on the X chromosome

Synonyms: ATLD2

Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: Base excision repair, Cell cycle, DNA replication, Mismatch repair, Nucleotide excision repair

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg Lane: Hela cells


Primary antibody: TA321511 (PCNA Antibody) at

dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA321511 (PCNA Antibody) at dilution 1/44 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human gasrtic cancer tissue using TA321511 (PCNA Antibody) at dilution 1/44, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA321511 (PCNA Antibody) at dilution 1/44 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using TA321511 (PCNA Antibody) at dilution 1/44, treated with fusion protein. (Original magnification: ×200)