

## **Product datasheet for TA321596S**

## **EDG6 (S1PR4) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Rat kidney tissue

IHC: 25-100

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide corresponding to a region derived from 352-367 amino acids of human

sphingosine-1-phosphate receptor 4

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 42 kDa

**Gene Name:** sphingosine-1-phosphate receptor 4

Database Link: NP 003766

Entrez Gene 13611 MouseEntrez Gene 8698 Human

O95977



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Background:** This gene is a member of the endothelial differentiation; G-protein-coupled (EDG)) receptor

gene family. EDG receptors bind lysophospholipids or lysosphingolipids as ligands; and are involved in cell signalling in many different cell types. This EDG receptor gene is intronless and is specifically expressed in the lymphoid tissue. Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse


physiological effect on most types of cells and tissues.

Synonyms: EDG6; LPC1; S1P4; SLP4

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 50 μg

Lane: Rat kidney tissue


Primary antibody: [TA321596] (S1PR4 Antibody)

at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution


Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321596] (S1PR4 Antibody) at dilution 1/40 (Original

magnification: ×200)





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using [TA321596] (S1PR4 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)