

# **Product datasheet for TA321602**

## **FADD Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Mouse spleen tissue and RAW264.7 cells

IHC: 50-100

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Full length fusion protein

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 23 kDa

**Gene Name:** Fas associated via death domain

Database Link: NP 003815

Entrez Gene 14082 MouseEntrez Gene 8772 Human

Q13158



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

#### **FADD Rabbit Polyclonal Antibody - TA321602**

Background:

The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain; this protein can be recruited by TNFRSF6/Fas-receptor; tumor necrosis factor receptor; TNFRSF25; and TNFSF10/TRAIL-receptor; and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein; which allows it to recruit caspase-8; and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.

Synonyms: GIG3; MORT1

**Protein Families:** Druggable Genome

**Protein Pathways:** Alzheimer's disease, Apoptosis, Pathways in cancer, RIG-I-like receptor signaling pathway,

Toll-like receptor signaling pathway

### **Product images:**



Gel: 8+12%SDS-PAGE Lysate: 30 µg

Lane 1-2: Mouse spleen tissue

RAW264.7 cells

Primary antibody: TA321602 (FADD Antibody) at


dilution 1/550

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 5 seconds






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321602 (FADD Antibody) at dilution 1/35 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321602 (FADD Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA321602 (FADD Antibody) at dilution 1/35 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA321602 (FADD Antibody) at dilution 1/35, treated with fusion protein. (Original magnification: ×200)