

Product datasheet for TA321608S

HDAC3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 200-1000

WB positive control: Raji cells

Reactivity: Human, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide corresponding to a region derived from 414-428 amino acids of Human

histone deacetylase 3

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Purification: Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49 kDa

Gene Name: histone deacetylase 3

Database Link: NP 003874

Entrez Gene 84578 RatEntrez Gene 8841 Human

<u>O15379</u>

Background: Histones play a critical role in transcriptional regulation; cell cycle progression; and

developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family. It has histone deacetylase activity and represses transcription when tethered to a promoter. It may participate in the regulation of

transcription through its binding with the zinc-finger transcription factor YY1. This protein can also down-regulate p53 function and thus modulate cell growth and apoptosis. This gene is

regarded as a potential tumor suppressor gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: HD3; RPD3; RPD3-2

Protein Families: Druggable Genome, Transcription Factors

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg Lane: Raji cells

Primary antibody: [TA321608] (HDAC3 Antibody)

at dilution 1/200

Secondary antibody: Goat anti rabbit IgG at

1/8000 dilution

Exposure time: 3 minutes