

## **Product datasheet for TA321661S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Transglutaminase 2 (TGM2) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Lovo and hela cells, human liver cancer tissue

IHC: 25-100

Positive control: Human cervical cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

Immunogen: Fusion protein corresponding to C terminal 250 amino acids of human transglutaminase 2 (C

polypeptide, protein-glutamine-gamma-glutamyltransferase)

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 77 kDa

**Gene Name:** transglutaminase 2

Database Link: NP 004604

Entrez Gene 21817 MouseEntrez Gene 7052 Human

P21980






Background:

Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved; they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. The protein encoded by this gene acts as a monomer; is induced by retinoic acid; and appears to be involved in apoptosis. Finally; the encoded protein is the autoantigen implicated in celiac disease. Two transcript variants encoding different isoforms have been found for this gene.

**Synonyms:** G-ALPHA-h; GNAH; HEL-S-45; TG2; TGC

Protein Families: Druggable Genome
Protein Pathways: Huntington's disease

## **Product images:**



Gel: 10%SDS-PAGE Lysate: 50 µg Lane 1-3: Lovo cells hela cells human liver cancer tissue


Primary antibody: [TA321661] (TGM2 Antibody) at

dilution 1/150

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute




Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321661] (TGM2 Antibody) at dilution 1/15 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA321661] (TGM2 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA321661] (TGM2 Antibody) at dilution 1/15 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colon cancer tissue using [TA321661] (TGM2 Antibody) at dilution 1/15, treated with fusion protein. (Original magnification: ×200)