OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for TA321722

RAD50 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	ELISA: 1:2000-10000, WB: 1:1000-5000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide corresponding to a region derived from 325-339 amino acids of Human RAD50 homolog (S. cerevisiae)
Formulation:	PBS pH7.3, 0.05% NaN3, 50% glycerol
Concentration:	lot specific
Purification:	Antigen affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	154 kDa
Gene Name:	RAD50 double strand break repair protein
Database Link:	NP_005723 Entrez Gene 19360 MouseEntrez Gene 64012 RatEntrez Gene 10111 Human Q92878
Background:	The protein encoded by this gene is highly similar to Saccharomyces cerevisiae Rad50; a protein involved in DNA double-strand break repair. This protein forms a complex with MRE11 and NBS1. The protein complex binds to DNA and displays numerous enzymatic activities that are required for nonhomologous joining of DNA ends. This protein; cooperating with its partners; is important for DNA double-strand break repair; cell cycle checkpoint activation; telomere maintenance; and meiotic recombination. Knockout studies of the mouse homolog suggest this gene is essential for cell growth and viability. Mutations in this gene are the cause of Nijmegen breakage syndrome-like disorder.

View online »

Synonyms: hRad50; NBSLD; RAD502

Protein Families: Druggable Genome

Protein Pathways: Homologous recombination, Non-homologous end-joining

Product images:



Predicted band size: 154 kDa. Positive control: 293T and Hela cell lysate. Recommended dilution: 1/1000-5000. (Gel: 8%SDS-PAGE Lane 1: 293T cell lysate Lane 2: Hela cell lysate Lysates: 40 ug per lane Primary antibody: 1/1500 dilution Secondary antibody: Goat anti Rabbit IgG - H&L (HRP) at 1/10000 dilution Exposure time: 5 minutes)