

## **Product datasheet for TA321735**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Melatonin Receptor 1A (MTNR1A) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: SP20 cell lysate

IHC: 30-150

Positive control: Human colorectal cancer Predicted cell location: Cell membrane

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide peptide corresponding to a region derived from 130-144 amino acids of

human melatonin receptor 1A

**Formulation:** PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 39 kDa

**Gene Name:** melatonin receptor 1A

Database Link: NP 005949

Entrez Gene 17773 MouseEntrez Gene 114211 RatEntrez Gene 4543 Human

P48039





Background:


This gene encodes one of two high affinity forms of a receptor for melatonin; the primary hormone secreted by the pineal gland. This receptor is a G-protein coupled; 7-transmembrane receptor that is responsible for melatonin effects on mammalian circadian rhythm and reproductive alterations affected by day length. The receptor is an integral membrane protein that is readily detectable and localized to two specific regions of the brain. The hypothalamic suprachiasmatic nucleus appears to be involved in circadian rhythm while the hypophysial pars tuberalis may be responsible for the reproductive effects of melatonin.

Synonyms: MEL-1A-R; MT1

**Protein Families:** Druggable Genome, GPCR, Transmembrane

**Protein Pathways:** Neuroactive ligand-receptor interaction

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: SP20 cell lysate


Primary antibody: TA321735 (MTNR1A Antibody)

at dilution 1/500

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 1 minute



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA321735 (MTNR1A Antibody) at dilution 1/25 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using TA321735 (MTNR1A Antibody) at dilution 1/25, treated with synthetic peptide. (Original magnification: ×200)