

Product datasheet for TA321746

ACAA2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: NIH/3T3 and A172 cells, human liver cancer tissue

IHC: 25-100

Positive control: Human colon cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Fusion protein corresponding to C terminal 250 amino acids of human acetyl-CoA Immunogen:

acyltransferase 2

Formulation: PBS pH7.3, 0.05% NaN3, 50% glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 42 kDa

Gene Name: acetyl-CoA acyltransferase 2

Database Link: NP 006102

Entrez Gene 52538 MouseEntrez Gene 170465 RatEntrez Gene 10449 Human

P42765



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background: 3-Ketoacyl-CoA thiolase, mitochondrial also known as acetyl-Coenzyme A acyltransferase 2 is

an enzyme that in humans is encoded by the ACAA2 gene. Acetyl-Coenzyme A

acyltransferase 2 is an acetyl-CoA C-acyltransferase enzyme. The encoded protein catalyzes the last step of the mitochondrial fatty acid beta-oxidation spiral. Unlike most mitochondrial


matrix proteins, it contains a non-cleavable amino-terminal targeting signal.

Synonyms: **DSAEC**

Protein Pathways: Fatty acid elongation in mitochondria, Fatty acid metabolism, Metabolic pathways, Valine,

leucine and isoleucine degradation

Product images:



Gel: 10%SDS-PAGE Lysate: 40 µg

Lane 1-3: NIH/3T3 cells

A172 cells

human liver cancer tissue


Primary antibody: TA321746 (ACAA2 Antibody) at

dilution 1/375

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 30 seconds




Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321746 (ACAA2 Antibody) at dilution 1/30 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human colon cancer tissue using TA321746 (ACAA2 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321746 (ACAA2 Antibody) at dilution 1/30 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using TA321746 (ACAA2 Antibody) at dilution 1/30, treated with fusion protein. (Original magnification: ×200)